

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssspta1204jxv

PASSWORD:

\* \* \* \* \* RECONNECTED TO STN INTERNATIONAL \* \* \* \* \*  
SESSION RESUMED IN FILE 'BEILSTEIN' AT 12:00:15 ON 02 MAY 2004  
FILE 'BEILSTEIN' ENTERED AT 12:00:15 ON 02 MAY 2004  
COPYRIGHT (c) 2004 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften  
licensed to Beilstein GmbH and MDL Information Systems GmbH

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	20.28	210.87

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-2.08

=> file reg

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	20.28	210.87

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-2.08

FILE 'REGISTRY' ENTERED AT 12:00:26 ON 02 MAY 2004  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
provided by InfoChem.

STRUCTURE FILE UPDATES: 30 APR 2004 HIGHEST RN 678535-01-8  
DICTIONARY FILE UPDATES: 30 APR 2004 HIGHEST RN 678535-01-8

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 6, 2004

Please note that search-term pricing does apply when  
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
information enter HELP PROP at an arrow prompt in the file or refer  
to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10231394.str


L6 STRUCTURE UPLOADED

=> d

L6 HAS NO ANSWERS

L6

STR



G1 C, Ch

Structure attributes must be viewed using STN Express query preparation.

=> s 16

SAMPLE SEARCH INITIATED 12:00:42 FILE 'REGISTRY'  
SAMPLE SCREEN SEARCH COMPLETED - 722 TO ITERATE

100.0% PROCESSED 722 ITERATIONS  
SEARCH TIME: 00.00.01

4 ANSWERS


FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*  
PROJECTED ITERATIONS: 12828 TO 16052  
PROJECTED ANSWERS: 4 TO 200

L7 4 SEA SSS SAM L6

=> d scan

L7 4 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfonyl]-, cis- (9CI)  
MF C16 H16 O4 S

Relative stereochemistry.




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):f10  
'F10' IS NOT VALID HERE

To display more answers, enter the number of answers you would like to see. To end the display, enter "NONE", "N", "0", or "END".  
HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):10


L7 4 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-1-methyl-2-(phenylsulfonyl)-, trans- (9CI)  
MF C16 H16 O3 S

Relative stereochemistry.




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L7 4 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 5-azido-2,3-dihydro-2-((3-methylphenyl)sulfinyl)-, acetate (ester)  
(9CI)  
MF C12 H13 N3 O4 S



L7 4 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-,  
[1α,2β(R\*)]- (9CI)  
MF C16 H16 O2 S

Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

ALL ANSWERS HAVE BEEN SCANNED

=> s 16 full  
FULL SEARCH INITIATED 12:01:37 FILE 'REGISTRY'  
FULL SCREEN SEARCH COMPLETED - 14799 TO ITERATE

100.0% PROCESSED 14799 ITERATIONS  
SEARCH TIME: 00.00.01

65 ANSWERS


L8 65 SEA SSS FUL L6

=> s l8 not sulfonyl  
955444 SULFONYL  
1 SULFONYLS  
955444 SULFONYL  
(SULFONYL OR SULFONYLS)

L9 30 L8 NOT SULFONYL

=> d scan

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 5-chloro-2,3-dihydro-2-(methylsulfinyl)-, acetate (9CI)  
MF C12 H13 Cl O3 S




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):10


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2S)-rel- (9CI)  
MF C15 H14 O2 S

Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-, acetate (9CI)  
MF C17 H15 Cl O3 S



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-,  
 [1α,2β(S\*)]- (9CI)  
 MF C16 H16 O2 S


Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1S-  
 [1α,2α(R\*)]]- (9CI)  
 MF C15 H14 O2 S


Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2,3-dihydro-1-methyl-2-(phenylsulfinyl)-,  
 (1α,2β)- (9CI)  
 MF C16 H16 O2 S

Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI)  
 MF C16 H16 O2 S



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-,  
 [1 $\alpha$ ,2 $\alpha$ (S\*)]- (9CI)  
 MF C16 H16 O2 S


Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-  
 [1 $\alpha$ ,2 $\beta$ (R\*)]]- (9CI)  
 MF C15 H14 O2 S

Absolute stereochemistry.




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN Benz[b]indeno[2,1-e]pyran-10(4bH)-one, 10a,11-dihydro-11-hydroxy-6,8-  
 dimethyl-10a-(methylsulfinyl)- (9CI)  
 MF C19 H18 O4 S



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1-Indanol, 2-[(p-butylphenyl)sulfinyl]- (7CI)  
 MF C19 H22 O2 S




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):10

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfinyl]-,  
 [1α,2β(R\*)]- (9CI)  
 MF C16 H16 O3 S


Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-,  
 [1α,2β(S\*)]- (9CI)  
 MF C15 H13 Cl O2 S


Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, acetate, (1R,2S)-rel-  
 (9CI)  
 MF C17 H16 O3 S


Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1-Indanol, 2-(dodecylsulfinyl)- (6CI)  
 MF C21 H34 O2 S





\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI)  
 MF C15 H13 Cl O2 S



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-,  
 [1 $\alpha$ ,2 $\alpha$ (S\*)]- (9CI)  
 MF C15 H13 Cl O2 S

Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 5-chloro-2,3-dihydro-2-(methylsulfinyl)- (9CI)  
 MF C10 H11 Cl O2 S



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2R)-rel- (9CI)  
MF C15 H14 O2 S

Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI)  
MF C15 H14 O2 S



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-, [1 $\alpha$ ,2 $\beta$ (R\*)]- (9CI)  
MF C16 H16 O2 S

Relative stereochemistry.




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):10

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1S-  
[1 $\alpha$ ,2 $\alpha$ (S\*)]]- (9CI)  
MF C15 H14 O2 S


Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*


L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-1-methyl-2-(phenylsulfinyl)-,  
(1 $\alpha$ ,2 $\alpha$ ) - (9CI)  
MF C16 H16 O2 S

Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI)  
MF C19 H16 O2 S




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfinyl]-,  
[1 $\alpha$ ,2 $\beta$ (S\*)] - (9CI)

MF C16 H16 O3 S


Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-  
[1 $\alpha$ ,2 $\beta$ (S\*)]]- (9CI)  
MF C15 H14 O2 S


Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, acetate, (1R,2R)-rel-  
(9CI)  
MF C17 H16 O3 S


Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
IN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1R,2R)-rel- (9CI)  
MF C15 H14 O2 S


Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfinyl]-,  
 [1α,2α(S\*)]- (9CI)  
 MF C16 H16 O3 S


Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

L9 30 ANSWERS REGISTRY COPYRIGHT 2004 ACS on STN  
 IN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-,  
 [1α,2β(R\*)]- (9CI)  
 MF C15 H13 Cl O2 S

Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

ALL ANSWERS HAVE BEEN SCANNED

=> file reg

COST IN U.S. DOLLARS

FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
160.69	371.56

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-2.08

FILE 'REGISTRY' ENTERED AT 12:02:30 ON 02 MAY 2004  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
 provided by InfoChem.

STRUCTURE FILE UPDATES: 30 APR 2004 HIGHEST RN 678535-01-8  
 DICTIONARY FILE UPDATES: 30 APR 2004 HIGHEST RN 678535-01-8

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 6, 2004

Please note that search-term pricing does apply when  
 conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more  
 information enter HELP PROP at an arrow prompt in the file or refer  
 to the file summary sheet on the web at:  
<http://www.cas.org/ONLINE/DBSS/registryss.html>

```
=> s 19
      955444 SULFONYL
        1 SULFONYLS
      955444 SULFONYL
          (SULFONYL OR SULFONYLS)
L10      30 L8 NOT SULFONYL
```

=> file caplus		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	4.85	376.41
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-2.08

FILE 'CAPLUS' ENTERED AT 12:02:37 ON 02 MAY 2004  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is  
 held by the publishers listed in the PUBLISHER (PB) field (available  
 for records published or updated in Chemical Abstracts after December  
 26, 1996), unless otherwise indicated in the original publications.  
 The CA Lexicon is the copyrighted intellectual property of the  
 American Chemical Society and is provided to assist you in searching  
 databases on STN. Any dissemination, distribution, copying, or storing  
 of this information, without the prior written consent of CAS, is  
 strictly prohibited.

FILE COVERS 1907 - 2 May 2004 VOL 140 ISS 19  
 FILE LAST UPDATED: 30 Apr 2004 (20040430/ED)

This file contains CAS Registry Numbers for easy and accurate  
 substance identification.

=> s 19

L11 22 L9

=> d hitstr 1-22

L11 ANSWER 1 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN


IT 500910-96-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; preparation of indanols as appetite depressants)

RN 500910-96-3 CAPLUS

CN 1H-Inden-1-ol, 5-chloro-2,3-dihydro-2-(methylsulfinyl)- (9CI) (CA INDEX NAME)



L11 ANSWER 2 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

IT 95720-00-6P 134779-81-0P 134779-82-1P

500910-96-3P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of indanols for producing drugs for prophylaxis or treatment of obesity)

RN 95720-00-6 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1R,2R)-rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 134779-81-0 CAPLUS


CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2S)-rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 134779-82-1 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2R)-rel- (9CI)  
 (CA INDEX NAME)

Relative stereochemistry.




RN 500910-96-3 CAPLUS  
 CN 1H-Inden-1-ol, 5-chloro-2,3-dihydro-2-(methylsulfinyl)- (9CI) (CA INDEX NAME)




L11 ANSWER 3 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 IT 500770-90-1 500770-91-2 500770-94-5  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (use of C2-substituted indan-1-ol derivs. in antiobesity drugs)  
 RN 500770-90-1 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, acetate, (1R,2R)-rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 500770-91-2 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, acetate, (1R,2S)-rel- (9CI) (CA INDEX NAME)

Relative stereochemistry.






RN 500770-94-5 CAPLUS  
 CN 1H-Inden-1-ol, 5-chloro-2,3-dihydro-2-(methylsulfinyl)-, acetate (9CI)  
 (CA INDEX NAME)




L11 ANSWER 4 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 IT 249514-81-6P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (preparation of 10,11-dihydro-11-hydroxybenz[b]indeno[2,1-e]pyran-10-ones  
 and analogs for enhancing biosynthesis of erythropoietin)  
 RN 249514-81-6 CAPLUS  
 CN Benz[b]indeno[2,1-e]pyran-10(4bH)-one, 10a,11-dihydro-11-hydroxy-6,8-  
 dimethyl-10a-(methylsulfinyl)- (9CI) (CA INDEX NAME)




L11 ANSWER 5 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 IT 155519-28-1P 155519-29-2P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)  
 RN 155519-28-1 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-1-methyl-2-(phenylsulfinyl)-,  
 (1 $\alpha$ ,2 $\beta$ )- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 155519-29-2 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-1-methyl-2-(phenylsulfinyl)-,  
 (1 $\alpha$ ,2 $\alpha$ )- (9CI) (CA INDEX NAME)

Relative stereochemistry.




L11 ANSWER 6 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
IT **32819-87-7P 32819-88-8P**  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 32819-87-7 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-  
[1α,2β(R\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 32819-88-8 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-  
[1α,2β(S\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 7 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
IT **92621-28-8P**  
RL: FORM (Formation, nonpreparative); PREP (Preparation)  
(formation of, in reaction of indene with thiophenol and oxygen, jet  
fuel instability model study in relation to)  
RN 92621-28-8 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)



L11 ANSWER 8 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN


IT 95720-00-6P 134779-81-0P 134779-82-1P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 95720-00-6 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1R,2R)-rel- (9CI)  
(CA INDEX NAME)


Relative stereochemistry.



RN 134779-81-0 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2S)-rel- (9CI)  
(CA INDEX NAME)


Relative stereochemistry.



RN 134779-82-1 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2R)-rel- (9CI)  
(CA INDEX NAME)

Relative stereochemistry.



L11 ANSWER 9 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN


IT 95720-00-6P

RL: FORM (Formation, nonpreparative); PREP (Preparation)  
(formation of, in indene reaction with thiophenol and oxygen or tert-Bu  
hydroperoxide, jet fuel in relation to)

RN 95720-00-6 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1R,2R)-rel- (9CI)  
(CA INDEX NAME)

Relative stereochemistry.



L11 ANSWER 10 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

IT **95720-00-6**


RL: PROC (Process)

(stereospecific formation of, in presence of ovalbumin)

RN 95720-00-6 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1R,2R)-rel- (9CI)  
(CA INDEX NAME)

Relative stereochemistry.



L11 ANSWER 11 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN


IT **62967-56-0**

RL: RCT (Reactant); RACT (Reactant or reagent)

(oxidation of, 1-indanone analog from)

RN 62967-56-0 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX NAME)




IT **65495-98-9P**

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 65495-98-9 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-, acetate (9CI)  
(CA INDEX NAME)




L11 ANSWER 12 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

IT 62967-56-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and Jones oxidation of)

RN 62967-56-0 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX NAME)



L11 ANSWER 13 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

IT 62703-00-8P 62703-01-9P 62703-02-0P

62703-05-3P 62703-06-4P 62703-07-5P


62703-10-0P 62703-11-1P 62703-12-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation and oxidation of)

RN 62703-00-8 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-,  
[1 $\alpha$ ,2 $\beta$ (S\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-01-9 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-,  
[1 $\alpha$ ,2 $\beta$ (R\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-02-0 CAPLUS


CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-,  
[1 $\alpha$ ,2 $\alpha$ (S\*)]- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 62703-05-3 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-,  
 [1α,2β(S\*)]- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 62703-06-4 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-,  
 [1α,2β(R\*)]- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 62703-07-5 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-,  
 [1α,2α(S\*)]- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 62703-10-0 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfinyl]-,  
 [1α,2β(S\*)]- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 62703-11-1 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfinyl]-,  
[1α,2β(R\*)]- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 62703-12-2 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfinyl]-,  
[1α,2α(S\*)]- (9CI) (CA INDEX NAME)

Relative stereochemistry.



L11 ANSWER 14 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
IT 32785-03-8P 32819-85-5P 32819-87-7P  
32819-88-8P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 32785-03-8 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1S-  
[1α,2α(R\*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 32819-85-5 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1S-

[1 $\alpha$ ,2 $\alpha$ (S\*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 32819-87-7 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-[1 $\alpha$ ,2 $\beta$ (R\*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 32819-88-8 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-[1 $\alpha$ ,2 $\beta$ (S\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L11 ANSWER 15 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

IT 62967-56-0, 1-Indanol, 2-[(p-chlorophenyl)sulfinyl]-  
92621-28-8, 1-Indanol, 2-(phenylsulfinyl)- 93434-18-5,  
1-Indanol, 2-(p-tolylsulfinyl)- 94305-49-4, 1-Indanol,  
2-(2-naphthylsulfinyl)- 102456-32-6, 1-Indanol,  
2-(dodecylsulfinyl)-  
(preparation of)


RN 62967-56-0 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX NAME)







RN 92621-28-8 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)




RN 93434-18-5 CAPLUS  
 CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 94305-49-4 CAPLUS  
 CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 102456-32-6 CAPLUS  
 CN 1-Indanol, 2-(dodecylsulfinyl)- (6CI) (CA INDEX NAME)




L11 ANSWER 16 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 IT 62967-56-0, 1-Indanol, 2-[(p-chlorophenyl)sulfinyl]-  
 92621-28-8, 1-Indanol, 2-(phenylsulfinyl)- 93434-18-5,  
 1-Indanol, 2-(p-tolylsulfinyl)- 94305-49-4, 1-Indanol,  
 2-(2-naphthylsulfinyl)- 94385-24-7, 1-Indanol,  
 2-[(p-butylphenyl)sulfinyl]- 102456-32-6, 1-Indanol,  
 2-(dodecylsulfinyl)-  
 (preparation of)  
 RN 62967-56-0 CAPLUS  
 CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX NAME)




RN 92621-28-8 CAPLUS  
 CN 1H-Inden-1-ol, 2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)




RN 93434-18-5 CAPLUS  
 CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 94305-49-4 CAPLUS  
 CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 94385-24-7 CAPLUS  
 CN 1-Indanol, 2-[(p-butylphenyl)sulfinyl]- (7CI) (CA INDEX NAME)




RN 102456-32-6 CAPLUS  
 CN 1-Indanol, 2-(dodecylsulfinyl)- (6CI) (CA INDEX NAME)




L11 ANSWER 17 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 IT **93434-18-5**, 1-Indanol, 2-(p-tolylsulfinyl)-  
 (preparation of)  
 RN 93434-18-5 CAPLUS  
 CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




L11 ANSWER 18 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 IT **93434-18-5**, 1-Indanol, 2-(p-tolylsulfinyl)-  
 (preparation of)  
 RN 93434-18-5 CAPLUS  
 CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




L11 ANSWER 19 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 IT **102456-32-6**, 1-Indanol, 2-(dodecylsulfinyl)-  
 (isomer)  
 RN 102456-32-6 CAPLUS  
 CN 1-Indanol, 2-(dodecylsulfinyl)- (6CI) (CA INDEX NAME)




IT **92621-28-8**, 1-Indanol, 2-(phenylsulfinyl)- **93434-18-5**,  
 1-Indanol, 2-p-tolylsulfinyl- **94305-49-4**, 1-Indanol,  
 2-(2-naphthylsulfinyl)-  
 (isomers)  
 RN 92621-28-8 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)




RN 93434-18-5 CAPLUS  
 CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 94305-49-4 CAPLUS  
 CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




IT 62967-56-0, 1-Indanol, 2-(p-chlorophenylsulfinyl)-  
 92621-28-8, 1-Indanol, 2-(phenylsulfinyl)- 93434-18-5,  
 1-Indanol, 2-p-tolylsulfinyl- 94305-49-4, 1-Indanol,  
 2-(2-naphthylsulfinyl)- 102456-32-6, 1-Indanol,  
 2-(dodecylsulfinyl)-  
 (preparation of)  
 RN 62967-56-0 CAPLUS  
 CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX NAME)




RN 92621-28-8 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)




RN 93434-18-5 CAPLUS  
 CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 94305-49-4 CAPLUS  
 CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 102456-32-6 CAPLUS  
 CN 1-Indanol, 2-(dodecylsulfinyl)- (6CI) (CA INDEX NAME)




L11 ANSWER 20 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 IT **92621-28-8**, 1-Indanol, 2-(phenylsulfinyl)-  
 (stereoisomers)  
 RN 92621-28-8 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)



L11 ANSWER 21 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
IT 94305-49-4, 1-Indanol, 2-(2-naphthylsulfinyl)-  
(preparation of)  
RN 94305-49-4 CAPLUS  
CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)



L11 ANSWER 22 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
IT 92621-28-8, 1-Indanol, 2-(phenylsulfinyl)-  
(stereoisomers)  
RN 92621-28-8 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)



=> s ibib abs hitstr 1-22  
14 IBIB  
212770 ABS  
5 ABSES  
212775 ABS  
(ABS OR ABSES)  
0 HITSTR  
7880518 1  
484522 22  
L12 0 IBIB ABS HITSTR 1-22  
(IBIB(W)ABS(W)HITSTR(W)1(W)22)

=> d ibib abs hitstr 1-22 l11

L11 ANSWER 1 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 2003:202618 CAPLUS  
DOCUMENT NUMBER: 138:221365  
TITLE: Preparation of indan-1-ols as appetite depressants  
INVENTOR(S): Jaehne, Gerhard; Krone, Volker; Bickel, Martin;  
Gossel, Matthias  
PATENT ASSIGNEE(S): Aventis Pharma Deutschland G.m.b.H., Germany  
SOURCE: PCT Int. Appl., 53 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: German  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----

WO 2003020696 A1 20030313 WO 2002-EP9206 20020817

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM  
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

DE 10142667 A1 20030327 DE 2001-10142667 20010831

US 2003114681 A1 20030619 US 2002-231394 20020830

US 6657086 B2 20031202


US 2004068016 A1 20040408 US 2003-665021 20030922

PRIORITY APPLN. INFO.: DE 2001-10142667 A 20010831


US 2002-231394 A3 20020830

OTHER SOURCE(S): MARPAT 138:221365

GI



I



II

AB Title compds. I [R1, R2, R3, R4 = H, halo, CN, etc.; X = S, SO, SO2; Y = (CH2)p; p = 0-3; R5 = CF3, alkyl, cycloalkyl] and their pharmaceutically acceptable salts were prepared For example, NaBH4 mediated reduction of 5-chloro-2-methylsulfonylindan-1-one, e.g., prepared from 2-bromo-5-chloroindan-1-one in 2-steps, provided indanol II. In milk consumption studies with female NMRI mice, indanol II exhibited very good anorectic effects, i.e., 50% decrease in milk consumption verses control.


IT 500910-96-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(drug candidate; preparation of indanols as appetite depressants)

RN 500910-96-3 CAPLUS

CN 1H-Inden-1-ol, 5-chloro-2,3-dihydro-2-(methylsulfinyl)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L11 ANSWER 2 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 2003:202465 CAPLUS

DOCUMENT NUMBER: 138:221361

TITLE: Preparation of indan-1-ols for producing drugs for the prophylaxis or treatment of obesity  
 INVENTOR(S): Jaehne, Gerhard; Krone, Volker; Bickel, Martin; Gossel, Matthias  
 PATENT ASSIGNEE(S): Aventis Pharma Deutschland G.m.b.H., Germany  
 SOURCE: PCT Int. Appl., 52 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003020263	A1	20030313	WO 2002-EP9205	20020817
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
DE 10142666	A1	20030320	DE 2001-10142666	20010831
US 2003134879	A1	20030717	US 2002-231183	20020830
PRIORITY APPLN. INFO.:			DE 2001-10142666 A	20010831
OTHER SOURCE(S):			MARPAT 138:221361	
GI				



AB Title compds. [I; R1-R4 = H, F, Cl, Br, I, cyano, N3, NO2, OH, alkoxy, cycloalkoxy, benzyloxy, phenoxy, alkylcarbonyloxy, etc.; X = S, SO, SO2; Y = (CH2)p; p = 0-3; R5 = CF3, (fluorinated) alkyl, cycloalkyl, etc.], were prepd for producing a drug for body weight loss of mammals. Thus, 5-chloro-2-methylsulfonylindan-1-one (preparation given) and NaBH4 in EtOH were put into a ultrasound bath for 4 h followed by stirring with 2N HCl to give 5-chloro-2-methylsulfonylindan-1-ol. The latter at 20 mg/kg i.p. was applied in female NMRI mice and gave 50% reduction of milk consumption of the treated mice.

IT 95720-00-6P 134779-81-0P 134779-82-1P  
 500910-96-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(preparation of indanols for producing drugs for prophylaxis or treatment of obesity)

RN 95720-00-6 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1R,2R)-rel- (9CI)  
 (CA INDEX NAME)


Relative stereochemistry.






RN 134779-81-0 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2S)-rel- (9CI)  
 (CA INDEX NAME)

Relative stereochemistry.




RN 134779-82-1 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2R)-rel- (9CI)  
 (CA INDEX NAME)

Relative stereochemistry.



RN 500910-96-3 CAPLUS  
 CN 1H-Inden-1-ol, 5-chloro-2,3-dihydro-2-(methylsulfinyl)- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 3 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 ACCESSION NUMBER: 2003:202409 CAPLUS  
 DOCUMENT NUMBER: 138:226750  
 TITLE: Use of C2-substituted indan-1-ol derivatives in antiobesity drugs  
 INVENTOR(S): Jaehne, Gerhard; Krone, Volker; Bickel, Martin; Gossel, Matthias

PATENT ASSIGNEE(S): Aventis Pharma Deutschland G.m.b.H., Germany  
 SOURCE: PCT Int. Appl., 54 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003020199	A1	20030313	WO 2002-EP9199	20020817
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
DE 10142660	A1	20030320	DE 2001-10142660	20010831
US 2003134881	A1	20030717	US 2002-230379	20020829
US 6667345	B2	20031223		

PRIORITY APPLN. INFO.: DE 2001-10142660 A 20010831  
 OTHER SOURCE(S): MARPAT 138:226750  
 GI



AB The invention relates to the use of C2-substituted indan-1-ol systems, and to the physiol. tolerable salts and the physiol. functional derivs. of the same, for producing medicaments used to reduce the weight of mammals, and for the prophylaxis or the treatment of obesity. The invention also relates to the use of compds. of formula (I), wherein the radicals have the cited designations, and to the physiol. tolerable salts and the physiol. functional derivs. of the same, for producing a medicament for the prophylaxis or the treatment of obesity. The antiobesity drugs can be combined with other active ingredients, e.g. cathine, phenylpropanolamine, amfepramone, mefenorex. Capsules, tablets, emulsions, dragees and suppositories are prepared containing the indan-1-ol derivative antiobesity drugs.


IT 500770-90-1 500770-91-2 500770-94-5

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (use of C2-substituted indan-1-ol derivs. in antiobesity drugs)

RN 500770-90-1 CAPLUS


CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, acetate, (1R,2R)-rel-  
 (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 500770-91-2 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, acetate, (1R,2S)-rel-  
 (9CI) (CA INDEX NAME)

Relative stereochemistry.



RN 500770-94-5 CAPLUS  
 CN 1H-Inden-1-ol, 5-chloro-2,3-dihydro-2-(methylsulfinyl)-, acetate (9CI)  
 (CA INDEX NAME)



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 4 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1999:733853 CAPLUS

DOCUMENT NUMBER: 131:322537

TITLE: Preparation of 10,11-dihydro-11-hydroxybenz[b]indeno[2,1-e]pyran-10-ones and analogs  
 for enhancing biosynthesis of erythropoietin

INVENTOR(S): Williams, Jonathan Gareth; Houck, David R.; Smith,  
 David Edward; Rathbone, Daniel Lee; Billington, David  
 Charles; Golding, Bernard T.; Collington, Eric W.;  
 Kitchin, John; Rich, Nicholas

PATENT ASSIGNEE(S): OSI Pharmaceuticals, Inc., USA

SOURCE: U.S., 13 pp., Cont.-in-part of U.S. 5,882,436.

CODEN: USXXAM

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5985913	A	19991116	US 1998-69693	19980429
PRIORITY APPLN. INFO.:			US 1996-32268P	P 19961129
			US 1997-978346	A2 19971126

OTHER SOURCE(S) : MARPAT 131:322537  
GI




AB Title compds. [I; R1,R4 = H or 1-4 of halo, alkyl, alkoxy, etc.; R2 = OR10 and R3 = H or R2R3 = O; R10 = H or alkanoyl; dashed line = optional addnl. bond] were prepared. Thus, 3',5'-dimethyl-2'-hydroxy-2-methylsulfinylacetophenone (preparation given) was cyclocondensed with 2-(OHC)C6H4CHO to give I (R1 = 6,8-Me2, R2 = OH, R3 = R4 = H, dashed line = bond). Data for biol. activity of I were given.

IT 249514-81-6P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(preparation of 10,11-dihydro-11-hydroxybenz[b]indeno[2,1-e]pyran-10-ones and analogs for enhancing biosynthesis of erythropoietin)

RN 249514-81-6 CAPLUS

CN Benz[b]indeno[2,1-e]pyran-10(4bH)-one, 10a,11-dihydro-11-hydroxy-6,8-dimethyl-10a-(methylsulfinyl)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L11 ANSWER 5 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1994:409577 CAPLUS

DOCUMENT NUMBER: 121:9577

TITLE: Reactions of  $\eta^2$ -(2-acylaryl-C,O)tetracarbonylmanganese(I) complexes with some vinyl sulfur compounds

AUTHOR(S): Cambie, Richard C.; Rutledge, Peter S.; Welch, David R.; Woodgate, Paul D.

CORPORATE SOURCE: Department of Chemistry, University of Auckland, Private Bag 92019, Auckland, N. Z.

SOURCE: Journal of Organometallic Chemistry (1994), 467(2), 237-44  
CODEN: JORCAI; ISSN: 0022-328X

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 121:9577

AB The thermally promoted reactions of some Ph and diterpenoid

$\eta^2$ -(2-acylaryl-C,O)tetracarbonylmanganese(I) complexes with Ph vinyl sulfone, Me vinyl sulfone, or Ph vinyl sulfoxide, have been investigated. The major products from the diterpenoid complexes arises from insertion followed by reductive demetalation; cyclopenta-annulation, when it occurs, is a minor process. Liberation of the metal-free adducts from their Mn-containing precursors requires treatment with either acid or photolysis-oxidation


IT 155519-28-1P 155519-29-2P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 155519-28-1 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-1-methyl-2-(phenylsulfinyl)-,  
(1 $\alpha$ ,2 $\beta$ )- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 155519-29-2 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-1-methyl-2-(phenylsulfinyl)-,  
(1 $\alpha$ ,2 $\alpha$ )- (9CI) (CA INDEX NAME)

Relative stereochemistry.



L11 ANSWER 6 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1991:558285 CAPLUS

DOCUMENT NUMBER: 115:158285

TITLE: Cooxidation reaction of indene and aromatic thiols in the presence of ovalbumin

AUTHOR(S): Freer, Juanita; Fuentealba, Cecilia; Gonzalez, Elizabeth; Baeza, Jaime

CORPORATE SOURCE: Dep. Quim., Univ. Concepcion, Concepcion, Chile

SOURCE: Phosphorus, Sulfur and Silicon and the Related Elements (1991), 61(1-2), 41-8

CODEN: PSSLEC; ISSN: 1042-6507

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 115:158285

AB The thiol olefin cooxidn. reaction (TOCO) between indene and aromatic thiols in presence of ovalbumin has been studied in hexane. While this reaction under normal conditions leads to the formation of 6 products, in the presence of OVA give stereospecifically only the trans-2-phenylmercapto-1-indanol derivative on the protein surface.

IT 32819-87-7P 32819-88-8P


RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of)

RN 32819-87-7 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-[1 $\alpha$ ,2 $\beta$ (R\*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 32819-88-8 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-[1 $\alpha$ ,2 $\beta$ (S\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 7 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1991:539316 CAPLUS

DOCUMENT NUMBER: 115:139316

TITLE: Fuel instability model studies: the liquid-phase cooxidation of thiols and indene by oxygen

AUTHOR(S): Morris, Robert E.; Mushrush, George W.

CORPORATE SOURCE: Nav. Technol. Cent. Saf. Survivabil., Nav. Res. Lab., Washington, DC, 20375, USA

SOURCE: Energy & Fuels (1991), 5(5), 744-8  
CODEN: ENFUEM; ISSN: 0887-0624

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Instability problems in middle distillate fuels were correlated with the presence of both active olefin species and heteroat. compds. such as thiols. The type of S compound rather than the total S concentration is the key to

fuel instability reactions. Low concns. of thiols will act as radical traps to inhibit autoxidn. When added to a fuel, thiols accelerated the rate of Oxidation without a commensurate increase in peroxidn. Evidence for the oxidative addition of thiols to olefins was observed by studying the

addition of thiophenol to indene in a model fuel during stressing in both a model system at 100-120° and in the jet fuel thermal oxidation test apparatus at 350°. Similarities and differences were found in the 2 systems, with the product distribution being temperature dependent. This could account, in part, for the differences in thiol influences on autoxidn. observed in model systems and in fuels.


IT 92621-28-8P

RL: FORM (Formation, nonpreparative); PREP (Preparation)


(formation of, in reaction of indene with thiophenol and oxygen, jet fuel instability model study in relation to)

RN 92621-28-8 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)




L11 ANSWER 8 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 1991:448966 CAPLUS  
DOCUMENT NUMBER: 115:48966  
TITLE: Cooxidation between thiophenol and cyclopentene  
AUTHOR(S): Freer, Juanita; Palma, Graciela; Fuentealba, Cecilia;  
Pena, Monica; Baeza, Jaime  
CORPORATE SOURCE: Dep. Quim., Univ. Concepcion, Concepcion, Chile  
SOURCE: Boletin de la Sociedad Chilena de Quimica (1991),  
36(1), 11-16  
CODEN: BOCQAX; ISSN: 0366-1644  
DOCUMENT TYPE: Journal  
LANGUAGE: Spanish  
GI




AB Treatment of thiophenol with cyclopentene in the presence of O2 gave adduct I (R = H) as well as cooxidn. products, i.e., cis- and trans-I (R = OH) and the sulfoxide derivs. The reaction of thiophenol with indene gave similar results.  
IT 95720-00-6P 134779-81-0P 134779-82-1P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)  
RN 95720-00-6 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1R,2R)-rel- (9CI)  
(CA INDEX NAME)

Relative stereochemistry.




RN 134779-81-0 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2S)-rel- (9CI)  
(CA INDEX NAME)

Relative stereochemistry.



RN 134779-82-1 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1S,2R)-rel- (9CI)  
(CA INDEX NAME)

Relative stereochemistry.



L11 ANSWER 9 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1988:224013 CAPLUS

DOCUMENT NUMBER: 108:224013

TITLE: Liquid-phase oxidation of thiophenol and indene by  
tert-butyl hydroperoxide and oxygen

AUTHOR(S): Mushruch, George W.; Watkins, John M.; Hazlett, Robert  
N.; Hardy, Dennis R.; Eaton, Harold G.

CORPORATE SOURCE: Nav. Res. Lab. Code 6180, Washington, DC, 20375-5000,  
USA

SOURCE: Fuel Science & Technology International (1988), 6(2),  
165-83

CODEN: FSCTEG; ISSN: 0884-3759

DOCUMENT TYPE: Journal

LANGUAGE: English

AB tert-Bu hydroperoxide (I) or O initiated the oxidation of thiophenol in the presence of indene was examined in C<sub>6</sub>H<sub>6</sub> at 120°. The reaction is kinetically complex, but it was possible to relate the product distribution to a few competing reactions. The product mixture was determined for several reaction time periods. The product slate was similar for all time periods, but yields of the individual components varied significantly with increasing reaction time. Gaseous products included isobutylene and a trace of CH<sub>4</sub>. The major product from I was tert-BuOH. The major product observed from thiophenol was Ph<sub>2</sub>S<sub>2</sub>. Addition products included the major product 2-phenylthiyl indan. Oxidation products included indanols, indanones, and the sulfoxide and sulfone of the major product 2-phenylthiyl indan. Solvent participation was noted by trace amts. of toluene.

IT 95720-00-6P


RL: FORM (Formation, nonpreparative); PREP (Preparation)  
(formation of, in indene reaction with thiophenol and oxygen or tert-Bu hydroperoxide, jet fuel in relation to)

RN 95720-00-6 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1R,2R)-rel- (9CI)  
(CA INDEX NAME)


Relative stereochemistry.





L11 ANSWER 10 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 ACCESSION NUMBER: 1985:148840 CAPLUS  
 DOCUMENT NUMBER: 102:148840  
 TITLE: Stereoselective oxidative addition of benzenethiol to indene in the presence of ovalbumin  
 AUTHOR(S): Baeza, Jaime; Freer, Juanita; Palma, Graciela  
 CORPORATE SOURCE: Dep. Quim., Univ. Concepcion, Concepcion, Chile  
 SOURCE: Monatshefte fuer Chemie (1984), 115(11), 1369-71  
 CODEN: MOCMB7; ISSN: 0026-9247  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 102:148840  
 AB The oxidative addition of PhSH to indene in the presence of ovalbumin produces only trans-anti-2-phenylsulfinyl-1-indanol. This reaction may be considered as a biomimetic model of detoxification of certain hydrocarbons by the liver.  
 IT 95720-00-6  
 RL: PROC (Process)  
 (stereospecific formation of, in presence of ovalbumin)  
 RN 95720-00-6 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-[(R)-phenylsulfinyl]-, (1R,2R)-rel- (9CI)  
 (CA INDEX NAME)

Relative stereochemistry.




L11 ANSWER 11 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
 ACCESSION NUMBER: 1978:190435 CAPLUS  
 DOCUMENT NUMBER: 88:190435  
 TITLE: Thiol-olefin cooxidation reaction. 6. A new convenient route to 1-substituted indenenes. Indenone as dienophile in Diels-Alder reactions  
 AUTHOR(S): Szmant, H. Harry; Nanjundiah, Raghunath  
 CORPORATE SOURCE: Dep. Chem. Chem. Eng., Univ. Detroit, Detroit, MI, USA  
 SOURCE: Journal of Organic Chemistry (1978), 43(9), 1835-7  
 CODEN: JOCEAH; ISSN: 0022-3263  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 88:190435  
 AB 2-(4-Chlorophenylsulfinyl)-1-indanone was decomposed in refluxing toluene to give indenone which was trapped by cyclopentadiene, hexachlorocyclopentadiene, and anthracene to give the resp. Diels-Alder adducts.  
 IT 62967-56-0  
 RL: RCT (Reactant); RACT (Reactant or reagent)

(oxidation of, 1-indanone analog from)

RN 62967-56-0 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX NAME)




IT 65495-98-9P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(preparation of)

RN 65495-98-9 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-, acetate (9CI)  
(CA INDEX NAME)



L11 ANSWER 12 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1977:422824 CAPLUS

DOCUMENT NUMBER: 87:22824

TITLE: A new route to 1,2-indanedione

AUTHOR(S): Szmant, H. Harry; Nanjundiah, Raghunath

CORPORATE SOURCE: Dep. Chem. Chem. Eng., Univ. Detroit, Detroit, MI, USA

SOURCE: Organic Preparations and Procedures International  
(1977), 9(1), 35-8


CODEN: OPPIAK; ISSN: 0030-4948

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 87:22824

GI



I

AB O was bubbled through a mixture of indene and p-ClC<sub>6</sub>H<sub>4</sub>SH in isooctane at room temperature and the resultant mixture of isomeric indanol sulfoxides (I;

X =


CHOH; R = H; R<sub>1</sub> = p-ClC<sub>6</sub>H<sub>4</sub>SO) was oxidized with Jones reagent to give indanone sulfoxide (I; X = CO; R and R<sub>1</sub> as before), which was refluxed with MeOH in the presence of iodine and the product (I; X = CO; R = R<sub>1</sub> = OMe) was deketalized with EtOH-H<sub>2</sub>SO<sub>4</sub> at reflux to give 1,2-indandione [I; X = CO; (RR<sub>1</sub>) = O].

IT 62967-56-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
(Reactant or reagent)  
(preparation and Jones oxidation of)

RN 62967-56-0 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX  
NAME)



L11 ANSWER 13 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1977:170438 CAPLUS

DOCUMENT NUMBER: 86:170438

TITLE: The thiol-olefin cooxidation (TOCO) reaction. IV.  
Temperature effects on product distribution in the  
TOCO reaction of indene and aromatic thiols

AUTHOR(S): Szmant, H. H.; Mata, A. J.; Namis, A. J.;  
Panthanickal, A. M.

CORPORATE SOURCE: Dep. Chem., Univ. Detroit, Detroit, MI, USA

SOURCE: Tetrahedron (1976), 32(22), 2665-80

CODEN: TETRAB; ISSN: 0040-4020

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The stereochem. of the TOCO reaction of indene with RC<sub>6</sub>H<sub>4</sub>SH (R = 4-Cl,  
4-MeO, 3-Me) is temperature dependent. Increasing amts. of cis addition  
products

are formed as the temperature is lowered to -23° and raised to  
.apprx.60°. The sensitivity of the temperature effect depends on the  
electronic character of the substituent in RC<sub>6</sub>H<sub>4</sub>SH. The effect of  
solvent, addition of cumyl hydroperoxide, K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>, galvinoxyl, PhNOCMe<sub>3</sub>,  
Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, and Na tetrathionate, and the kinetics of the TOCO reaction were  
determined

IT 62703-00-8P 62703-01-9P 62703-02-0P

62703-05-3P 62703-06-4P 62703-07-5P


62703-10-0P 62703-11-1P 62703-12-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
(Reactant or reagent)  
(preparation and oxidation of)

RN 62703-00-8 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-,  
[1 $\alpha$ ,2 $\beta$ (S\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-01-9 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-,  
[1 $\alpha$ ,2 $\beta$ (R\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-02-0 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro-,  
[1α,2α(S\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-05-3 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-,  
[1α,2β(S\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-06-4 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-,  
[1α,2β(R\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-07-5 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(3-methylphenyl)sulfinyl]-,  
[1α,2α(S\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-10-0 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfinyl]-,  
[1α,2β(S\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-11-1 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfinyl]-,  
[1α,2β(R\*)]- (9CI) (CA INDEX NAME)


Relative stereochemistry.



RN 62703-12-2 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-[(4-methoxyphenyl)sulfinyl]-,  
[1α,2α(S\*)]- (9CI) (CA INDEX NAME)

Relative stereochemistry.



L11 ANSWER 14 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1972:85045 CAPLUS

DOCUMENT NUMBER: 76:85045

TITLE: Nonstereospecific oxidative addition of benzenethiol  
to indene

AUTHOR(S): Szmant, H. Harry; Rigau, Juan J.  
 CORPORATE SOURCE: Puerto Rico Nucl. Cent., Univ. Puerto Rico, San Juan, P. R.  
 SOURCE: Journal of Organic Chemistry (1972), 37(3), 447-51  
 CODEN: JOCEAH; ISSN: 0022-3263  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB The oxidative addition of benzenethiol to indene produces a mixture of three isomeric 2-phenylsulfinylindanol that contains 14-18% of the cis-hydroxy sulfoxide contrary to previous claims that this reaction leads stereospecifically only to trans addition products. The fourth isomeric 2-phenylsulfinylindanol (cis-syn) was prepared by the oxidation of the sulfide precursor. Examination of the concentration dependence of H bonding, of the

NMR spectra, and of the relative yields permits assignment of the configurations at the S atom in all isomers.


IT 32785-03-8P 32819-85-5P 32819-87-7P  
 32819-88-8P

RL: SPN (Synthetic preparation); PREP (Preparation)  
 (preparation of)

RN 32785-03-8 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1S-[1 $\alpha$ ,2 $\alpha$ (R\*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 32819-85-5 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1S-[1 $\alpha$ ,2 $\alpha$ (S\*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 32819-87-7 CAPLUS


CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-[1 $\alpha$ ,2 $\beta$ (R\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 32819-88-8 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)-, [1R-  
[1 $\alpha$ ,2 $\beta$ (S\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L11 ANSWER 15 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 1962:475774 CAPLUS  
DOCUMENT NUMBER: 57:75774  
ORIGINAL REFERENCE NO.: 57:15030e-i  
TITLE: Hydroperoxides and sulfoxides  
INVENTOR(S): Oswald, Alexis A.; Rupar, Charles B.; Greenwood,  
Sydney H.J.  
PATENT ASSIGNEE(S): Esso Research and Engineering Co.  
SOURCE: 8 pp.  
DOCUMENT TYPE: Patent  
LANGUAGE: Unavailable  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 3043824		19620710	US	19590928


AB Co-oxidation of RSH with olefins or diolefins in the presence of O at 0° produced new sulfoxides and hydroperoxides. Ultraviolet light and hydroperoxides were found to be effective catalysts for the oxidation. Thus, 20.2 g. n-C<sub>12</sub>H<sub>25</sub>SH and 10.4 g. styrene were dissolved in 150 cc. n-C<sub>7</sub>H<sub>16</sub> in 500 cc. quartz bottle and irradiated with ultraviolet light (or hydroperoxides added). Air was passed through the solution via sintered glass and filtered after 0.5 hr. to give 18.7% 2-phenyl-2-hydroperoxyethyl n-dodecyl sulfide. Similarly prepared were 1-hydroperoxy-4-phenylthio-2-butene, 1-hydroperoxy-4-(2-naphthylthio)-2-pentene, 2-hydroperoxy-4-(2-naphthylthio)-2cyclopentene, and 4,7-methylene-5-(2-naphthylthio)-6-hydroperoxy-4,5,6,7,8,9-hexahydroindene. The filtrate was treated with air 6 hrs. and filtered to give 9 g. 2-phenyl-2hydroxyethyl n-dodecyl sulfoxide (I), m. 107-9° (benzene-nheptane). The filtrate gave a mixture of isomeric I, m. 66-9°. Similarly prepared were the following sulfoxides, where R<sub>1</sub> = C<sub>16</sub>H<sub>33</sub>CH(OH)CH<sub>2</sub>, R<sub>2</sub> = p-MeC<sub>6</sub>H<sub>4</sub>, R<sub>3</sub> = naphthyl, R<sub>4</sub> = p-ClC<sub>6</sub>H<sub>4</sub>, R<sub>5</sub> = PhCH(OH)CH<sub>2</sub>, R<sub>6</sub> = PhMe(OH)CH<sub>2</sub>, R<sub>7</sub> = C<sub>12</sub>H<sub>25</sub>, and R<sub>8</sub> = 2-(1-hydroxyindanyl): PhSOCH<sub>2</sub>CH(OH)C<sub>10</sub>H<sub>21</sub>, m. 67-8°; PhSOR<sub>1</sub>, m. 51-2°; R<sub>2</sub>SOR<sub>1</sub>, m. 60-2°; R<sub>3</sub>SOR<sub>1</sub>, m. 88-90°; R<sub>4</sub>SOR<sub>1</sub>, m. 65-6.5°; R<sub>5</sub>SOPh, m. 123-30°; R<sub>5</sub>SOR<sub>3</sub>, m. 145-7°; R<sub>5</sub>SOR<sub>4</sub>, m. 156.5-8.5°; R<sub>6</sub>SOR<sub>3</sub>, m. 114-18°; R<sub>6</sub>SOR<sub>7</sub>, m. 47.5-8.5°; R<sub>8</sub>SOPh, m. 158-9° (decomposition); R<sub>8</sub>SOR<sub>2</sub>, 166-7.5° (decomposition); R<sub>8</sub>SOR<sub>4</sub> m. 144-6°; R<sub>8</sub>SOR<sub>3</sub>, m. 149-50°; and R<sub>8</sub>SOR<sub>7</sub>, m. 107-9°. The hydroxy sulfoxide products of dicyclopentadiene-RSH co-oxidns. were useful as petroleum additives, antistatic agents, and pesticides. The OH group may be esterified with H<sub>2</sub>SO<sub>4</sub> and converted to detergents. The hydroperoxides are useful as radical polymerization promoters and can be alkylated to give surfactive agents. Co-oxidation can be utilized to remove dienes from steam-cracked naphtha and for removal of RSH from petroleum.

IT 62967-56-0, 1-Indanol, 2-[(p-chlorophenyl)sulfinyl]-  
92621-28-8, 1-Indanol, 2-(phenylsulfinyl)- 93434-18-5,  
1-Indanol, 2-(p-tolylsulfinyl)- 94305-49-4, 1-Indanol,

2-(2-naphthylsulfinyl)- 102456-32-6, 1-Indanol,  
2-(dodecylsulfinyl)-  
(preparation of)


RN 62967-56-0 CAPLUS

CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX NAME)




RN 92621-28-8 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)




RN 93434-18-5 CAPLUS

CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 94305-49-4 CAPLUS

CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)



RN 102456-32-6 CAPLUS

CN 1-Indanol, 2-(dodecylsulfinyl)- (6CI) (CA INDEX NAME)





ACCESSION NUMBER: 1962:73337 CAPLUS  
DOCUMENT NUMBER: 56:73337  
ORIGINAL REFERENCE NO.: 56:14179h-i,14180a-i,14181a-b  
TITLE: Organic sulfur compounds. VI. The effect of  
alkylamines on the course of the cooxidation of  
mercaptans and indene  
AUTHOR(S): Oswald, Alexis A.; Noel, Fernand; Fisk, George  
CORPORATE SOURCE: Imp. Oil Co., Sarnia, Can.  
SOURCE: Journal of Organic Chemistry (1961), 26, 3974-80  
CODEN: JOCEAH; ISSN: 0022-3263  
DOCUMENT TYPE: Journal  
LANGUAGE: Unavailable

AB In the presence of alkylamines, mercaptans, and indene (I) were cooxidized by mol. O with a chain mechanism to form substituted 2-mercapto-1-indanols, disulfides, and H<sub>2</sub>O, instead of substituted 2-mercapto-1-indanyl hydroperoxides. The initiation reaction forming the mercapto radicals could take place between alkylammonium thiolates and O. The change of the reaction products was due to the catalysis by the amines of the oxidation of mercaptans by substituted 2-mercapto-1-indanyl hydroperoxides. It was proposed that this catalytic action was important in the stabilization of some hydrocarbon fuels by alkylamines. In the cooxidn. expts., the O or air was introduced through a sintered glass inductor, mixture stirred, and the pressure kept slightly above atmospheric. The reactions were followed by determining the decrease of the thiol concentration by potentiometric titration of samples with AgNO<sub>3</sub>. 2-(2-Naphthylthio)-1-indanyl hydroperoxide (II) (14.2 g.) in 2.5 l. PhMe, 6.6 g. benzenethiol (IIa), and 7.6 g. 1,1,3,3-tetramethylbutylamine (III) left 0.5 hrs., the solution evaporated in vacuo, at first 2-(2-naphthylsulfinyl)-1-indanol removed, then the solvent evaporated, and from the residue III and diphenyl disulfide removed by leaching with 2 60-ml. portions of heptane, and the product crystallized gave 5.2 g. 2-(2-naphthylthio)-1-indanol (IIIa), m. 141-3° (PhMe). IIa (13.2 g.) and 7.7 g. III in 250 ml. C<sub>6</sub>H<sub>6</sub> treated with 19.4 g. 79% 2-phenylthio-1-indanyl hydroperoxide, the mixture stirred another hr. and evaporated, and the residue crystallized gave 5 g. 2-phenylthio-1-indanol (IIIB), m. 103-4° (heptane). A yield of 4.2 g. (PhS)<sub>2</sub> was also obtained in this experiment II (6.3 g.) in 25 ml. Et<sub>2</sub>O added slowly to 1.1 g. triethylenediamine in 10 ml. Et<sub>2</sub>O (an exothermic reaction occurred) and the solid collected gave 6.7 g. triethylenediammonium 2-(2-naphthylthio)-1-indanyl peroxide (IV), m. 89-90°. The synthesis of IV was also accomplished in PhMe. IIa (1.1 g.) in 80 ml. PhMe treated with 1.8 g. IV (slowly), the mixture left 0.5 hr. and filtered, and the solid recrystd. gave 1.2 g. IIIa. 2-Naphthalenethiol (6.4 g.) in 50 ml. PhMe treated 0.5 hr. at room temperature with 7.3 g. IV gave 8.8 g. addition product (V) of 2-(2-naphthylthio)-1-indanol and di-2-naphthyl disulfide, m. 126-7° (PhMe). IIIa (1.46 g.) and 1.6 g. di-2-naphthyl disulfide in 25 ml. PhMe heated, cooled to room temperature, and the crystals collected gave 2.5 g. V. A C<sub>6</sub>H<sub>6</sub> solution (333 ml.) of 0.34 mole aromatic thiol, 0.11 mole I, and 0.01 mole of an alkylamine was oxygenated 6 hrs. at 22-8°. When benzenethiol and 2-naphthalenethiol were cooxidized with I in the presence of III, a yellow color developed. With 4-butylbenzenethiol, the mixture became dark yellow; with 4-toluenethiol and 4-chlorobenzenethiol, it became red and black, resp. When benzenethiol and I were cooxidized in the presence of mono-, di-, and tripropylamines, the formation of purple, green, and yellow colors were observed. After oxygenation, the C<sub>6</sub>H<sub>6</sub> layer was decanted, the unchanged thiol removed from the C<sub>6</sub>H<sub>6</sub> solution by washing with 5% KOH solution, the solvent distilled, and the residue fractionated to give 2-arylthio-1-indanols. The diaryl disulfides were too soluble in heptane and therefore were crystallized from MeOH. In the case of the 2-naphthalenethiol-I cooxidn., the reactant concns. were reduced to one-third, and C<sub>6</sub>H<sub>6</sub> used as solvent of recrystn. because of the solubility of

the addition compound of IIIa and di-2-naphthyl disulfide. The cooxidn. of IIa and I in the presence of III was also carried out on a 5 times larger scale in PhMe. After 6 hrs. of oxygenation, 58% of the thiol was oxidized and 4.4 g. of H<sub>2</sub>O separated. By the extraction of the reaction mixture with 5%

HCl

and the concentration of the extract, 7 g. III.HCl was isolated. Workup of the PhMe

solution gave 20.5 g. IIb. A heptane solution (333 ml.) of 73.3 g. dodecanethiol, 22.9 g. I, and 4.2 g. III was oxygenated at 24-6° and the product, m. 69-72°, isolated. In addition to the above, the following substituted 2-mercapto-1-indanols were prepared (substituent, m.p., and % yield given): 4-tolyl, 95.5-6.5°, 64; 4-butylphenyl, 104-5°, 39; 4-chlorophenyl, 113.5-14.5°, 35. 2-Arylthio-1-indanol (0.01 mole) in AcOH was treated slowly at 50° with 1.1 g. 33% H<sub>2</sub>O<sub>2</sub>, the mixture kept 0.5 hr. at 60°, the resulting 2-arylsulfinyl-1-indanol precipitated by addition of H<sub>2</sub>O, the solid collected,

and

recrystd. from alc. Most of the oxidation products were identified with one isomer of the substituted 2-sulfinyl-1-indanols from the cooxidn. of I with the corresponding thiol in the absence of amine as shown (thiol starting material, m.p. of the 2-sulfinyl-1-indanol obtained by H<sub>2</sub>O<sub>2</sub> oxidation given): benzene, 147-8.5° (decomposition); toluene, 144-5.5° (decomposition); 4-chlorobenzene, 146.5-8.0° (decomposition); 2-naphthalene, 134.5-6.0°; dodecane, 80-1°. An AcOH solution of 0.01 mole of 2-arylthio-1-indanol was oxidized with 2.3 g. 33% H<sub>2</sub>O<sub>2</sub> as described above; the work up of the mixture (after being heated 1 hr. at 80°) yielded the 2-arylsulfonyl-1-indanols given below. The following substituted 2-sulfinyl- and 2-sulfonylindanols were thus obtained (R and x of RSO<sub>x</sub>, m.p., and % yield given): 4-butylphenyl, 1, 154-6°, 63; 4-dodecyl, 1, 80-1°, 71; 4-tolyl, 2, 145-6.5°, 72; 4-butylphenyl, 2, 156-8°, 98; 4-chlorophenyl, 2, 151.5-4.0°, 95; dodecyl, 2, 93.5-4.5°, 97. Benzene solns. containing 0.15 mole/l. of benzenethiol and 0.05 mole/l. I were oxygenated in the presence of various concns. of III at room temperature. After 6 hrs. of oxygenation, the following decrease in % mercaptan was observed (amine mole/l., % thiol oxidized given): nil, 41; 0.015, 14; 0.150, 48; 0.300, 47. Primene 81-R was added to heptane solns. containing benzenethiol, I, and 2,5-dimethylpyrrole, the test solns. (300 ml. each) were aerated 6 hrs. at room temperature, and the following observations were made (mole/l.

each

of benzenethiol, I, and 2,5-dimethylpyrrole, mole/l. Primene 81-R, % thiol oxidized, peroxide formed, color of solution, and precipitate in g./100 ml.

given):

0.30, nil, 68, yes, red, 3.5 (red oil); 0.30, 0.03, 56, no, colorless, 1.3 (colorless crystals); 0.01, nil, 75, yes, red, 0.2 (red solid); 0.01, 0.001, 53, no, colorless, none.

IT


62967-56-0, 1-Indanol, 2-[(p-chlorophenyl)sulfinyl]-  
92621-28-8, 1-Indanol, 2-(phenylsulfinyl)- 93434-18-5,  
1-Indanol, 2-(p-tolylsulfinyl)- 94305-49-4, 1-Indanol,  
2-(2-naphthylsulfinyl)- 94385-24-7, 1-Indanol,  
2-[(p-butylphenyl)sulfinyl]- 102456-32-6, 1-Indanol,  
2-(dodecylsulfinyl)-  
(preparation of)

RN


62967-56-0 CAPLUS

CN


1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX NAME)




RN 92621-28-8 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)




RN 93434-18-5 CAPLUS  
CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 94305-49-4 CAPLUS  
CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)



RN 94385-24-7 CAPLUS  
CN 1-Indanol, 2-[(p-butylphenyl)sulfinyl]- (7CI) (CA INDEX NAME)




RN 102456-32-6 CAPLUS  
CN 1-Indanol, 2-(dodecylsulfinyl)- (6CI) (CA INDEX NAME)



ACCESSION NUMBER: 1962:47254 CAPLUS  
 DOCUMENT NUMBER: 56:47254  
 ORIGINAL REFERENCE NO.: 56:9005i,9006a  
 TITLE: Hydrocarbon purification process and catalyst  
 INVENTOR(S): Holden, Donald L.  
 PATENT ASSIGNEE(S): Universal Oil Products Co.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 PATENT INFORMATION:


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	US 3016348		19620109	US	19591102
AB	A catalyst is prepared by combining an Al <sub>2</sub> O <sub>3</sub> -SiO <sub>2</sub> carrier material containing 10-25% SiO <sub>2</sub> with 5-10% Mo and 1-5% Ni, then sulfiding so that the Mo and Ni exist as sulfides in the final composition. Comparison of the liquid hydrogenation products obtained by use of this catalyst with those obtained by use of the well-known Al-Co-Mo catalyst showed marked improvement in the removal of N compds.				
IT	93434-18-5, 1-Indanol, 2-(p-tolylsulfinyl)- (preparation of)				
RN	93434-18-5 CAPLUS				
CN	1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)				



L11 ANSWER 18 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1962:47253 CAPLUS  
 DOCUMENT NUMBER: 56:47253  
 ORIGINAL REFERENCE NO.: 56:9005h-i  
 TITLE: Denitrogenation of petroleum  
 INVENTOR(S): Oswald, Alexis A.; Rupar, Charles B.; Ilnyekyj, Stephen; Noel, Fernand  
 PATENT ASSIGNEE(S): Esso Research and Engineering Co.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 PATENT INFORMATION:

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	US 3016349		19620109	US	19590810
AB	Addition of 0.5 to 3 moles of an organic peroxide such as Bz <sub>2</sub> O <sub>2</sub> , (4-ClC <sub>6</sub> H <sub>4</sub> ) <sub>2</sub> O <sub>2</sub> , or tert-Bu <sub>2</sub> O <sub>2</sub> per mole of pyrroles in the petroleum product at 50-250°F. converts the pyrrole-type compds. to higher-boiling compds. which are separated by distillation or filtration. Sulfides, such as 2-indanyl or p-tolyl sulfide, or sulfoxides, such as 2-phenyl-2-hydroxyethyl phenyl sulfoxide, may also be used, but at 60-360°F.				
IT	93434-18-5, 1-Indanol, 2-(p-tolylsulfinyl)- (preparation of)				
RN	93434-18-5 CAPLUS				
CN	1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)				



L11 ANSWER 19 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1961:124710 CAPLUS

DOCUMENT NUMBER: 55:124710

ORIGINAL REFERENCE NO.: 55:23460b-i,23461a-c

TITLE: Organic sulfur compounds. III. Cooxidation of mercaptans with styrenes and indene

AUTHOR(S): Oswald, Alexis A.

CORPORATE SOURCE: Imp. Oil Ltd., Sarnia, Can.

SOURCE: Journal of Organic Chemistry (1961), 26, 842-6

CODEN: JOCEAH; ISSN: 0022-3263

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable


GI For diagram(s), see printed CA Issue.

AB cf. CA 54, 21005e. PhSH (I) (0.1 mole) and 0.1 mole PhCH:CH<sub>2</sub> (II) [or PhCMe:CH<sub>2</sub> (III) or indene (IV)] in 310 ml. cold n-C<sub>7</sub>H<sub>16</sub> bubbled through with air via a sintered glass sparger 6 hrs. at 0° yielded 30, 52, and 33% oily peroxides containing 84, 44, and 67% hydroperoxide, according to both the iodide and FeSO<sub>4</sub> methods. Similar aeration of p-ClC<sub>6</sub>H<sub>4</sub>SH and III yielded 64% peroxide containing 58% hydroperoxide. A 4-g. sample of liquid substituted 2-mercaptoethyl hydroperoxide kept 3 days at 20° and the solid product recrystd. from PhMe-C<sub>7</sub>H<sub>16</sub> yielded the corresponding isomeric 2-sulfinylethanol rearrangement products, RSO-CH<sub>2</sub>CR'PhOH (V). A typical member of these new hydroperoxides was 1-(2-naphthylthio)-2-phenyl-2-propyl hydroperoxide (VI), m. -10°, peroxide content 85%, obtained by cooxidn. of 2-HSC<sub>10</sub>H<sub>7</sub> and III. Phys. data were listed for V [R, R', m.p. (uncor.), and infrared OH, SO (2), aromatic and CH<sub>2</sub> absorption peaks given]: Ph, H, 128-9.5°, 2.95, 9.23 (9.55), 10.02, 6.3, 6.9; 2-Cl<sub>10</sub>H<sub>7</sub>, H, 145-7° (121-6°), 3 (3.05), 9.5 (9.45), 9.9 (9.9), 6.28 (6.28), 6.91 (6.91); Cl<sub>12</sub>H<sub>25</sub>, H, 105-6° (69-70°), 3 (3), 9.3 (9.3), 9.85 (9.85), 6.25 (6.25), 6.9 (6.9); 4-ClC<sub>6</sub>H<sub>4</sub>, H, 156.5-8.5° (86-8°), 2.96 (2.96), 9.19 (9.19), 9.94 (9.96), 6.35 (6.35), 6.95 (6.95); 2-Cl<sub>10</sub>H<sub>7</sub>, Me, 115-18° (94-95.5°), 3.05 (3), 9.4 (9.4), 9.95 (9.78), 6.28 (6.28), 6.85 (6.93); Cl<sub>12</sub>H<sub>25</sub>, Me, 47.5-8.5°, 2.95, 9.4, 10.05, 6.25, 6.85. Similar rearrangements were carried out in 0.3M C<sub>6</sub>H<sub>6</sub> (CHCl<sub>3</sub>, tetrahydronaphthalene) solns. of hydroperoxides at 43° in 16 hrs. VI (3.1 g.) yielded 27% V (R = 2-Cl<sub>10</sub>H<sub>7</sub>, R' = H), m. 115-18°, by this method. IV (11.6 g.) and 16 g. 2-Cl<sub>10</sub>H<sub>7</sub>SH in 100 ml. C<sub>6</sub>H<sub>6</sub> and 300 ml. C<sub>7</sub>H<sub>16</sub> aerated 4 hrs. at 0° yielded 4 g. crystalline 2-(2-naphthylthio)-1-indanyl hydroperoxide (VII), m. 70° (decomposition), containing 98% peroxide. The filtrate aerated 6 hrs. at 0° and 42 hrs. at 20°, filtered, and the cooxidn. product (26 g., containing only 2% peroxide) recrystd. from C<sub>6</sub>H<sub>6</sub>-C<sub>7</sub>H<sub>16</sub> gave the 2-(2-naphthylsulfinyl)-2-indanoisomers, C<sub>6</sub>H<sub>4</sub>.CH(OH).CH(SOR).CH<sub>2</sub> (VIII, R = 2-Cl<sub>10</sub>H<sub>7</sub>) (IX), m. 157-8°, m. 149-50°, m. 138.5-9.5° (decomposition), m. 125-7.5° (decomposition) (infrared spectrum given for all). VII (9.2 g.) in 100 ml. C<sub>6</sub>H<sub>6</sub> kept 16 hrs. at 43° gave a neg. peroxide test and deposited 36.8 g. crystals, m. 138.5-9.5° (C<sub>6</sub>H<sub>6</sub>), identical with an isomer of IX. VII stored 10 min. in a desiccator at 2 mm. over H<sub>2</sub>SO<sub>4</sub> decomposed violently and the reddish product fractionally crystallized from


C<sub>6</sub>H<sub>6</sub> gave IX, m. 157-8° and 138.5-9.5°. IV (10.7 g.) and 12.8 g. 4-ClC<sub>6</sub>H<sub>4</sub>SH in 250 ml. C<sub>6</sub>H<sub>6</sub> and 20 ml. PhCl at 0° aerated 4 hrs. and the fresh peroxide solution refluxed 1 hr. in Et<sub>2</sub>O with LiAlH<sub>4</sub> yielded 68% 2-(4-chlorophenylthio)-1-indanol (X), m. 110-12°. VII

(6.0 g.) in 600 ml. C<sub>6</sub>H<sub>6</sub>, 150 ml. CHCl<sub>3</sub>, and 20 ml. MeOH kept 3 days at 20° with 4.3 g. X gave unchanged X and IX, m. 157-8°, suggesting an intramol. rearrangement. However, the reactions of the hydroperoxide cooxidn. intermediate might involve radical intermediates as was suggested by the polymerization of II by VII. VII (0.3 g.) in 34.6 g. II kept 3 months at 5° gave only 75% recovered II on distillation at 100°/20 mm., whereas II kept alone under the same conditions was distilled with 99% recovery. Aliphatic hydrocarbon (C<sub>7</sub>H<sub>16</sub>, cetane, straight run petroleum distillate, b. 70-200°) (300 ml.) containing 0.1 mole thiol, p-RC<sub>6</sub>H<sub>4</sub>SH (R = H, Me, 4-ClC<sub>6</sub>H<sub>4</sub>, and II, III, or IV aerated (or oxygenated) 3 days at 20° to give a liquid phase with peroxide number and mercaptan number less than 5 and 10, resp., the residue on filtration recrystd. from Ph-Me-C<sub>7</sub>H<sub>16</sub> to give 40-80% mixts. of isomeric substituted 2-sulfinyl ethanols, and the mixts. recrystd. gave (with about 30% yield loss) the isomeric VIII (R and m.p. given) (infrared spectra given): Ph, 158-9°; 148-50°, 132-5°; p-MeC<sub>6</sub>H<sub>4</sub>, 166-7.5°, 144-4.5°, 128-30°; p-ClC<sub>6</sub>H<sub>4</sub>, 144-6°; C<sub>12</sub>H<sub>25</sub>, 107-9°, 67-8.5°. C<sub>12</sub>H<sub>25</sub>SH (20.2 g.) and 0.1 mole II (III, IV) in 300 ml. C<sub>7</sub>H<sub>16</sub> aerated in a Vycor (95% quartz) flask with ultraviolet irradiation (GE-9T64Y20 lamp, 250 v., 1000 w. at 4.5 cm.) with decrease to less than 10% thiol content, the mixture cooled (solid CO<sub>2</sub>-alc. bath), and the product recryst. gave 75% VIII (R = C<sub>12</sub>H<sub>25</sub>) isomers. Cooxidn. of C<sub>12</sub>H<sub>25</sub>SH with III and IV gave only 10% V (R = C<sub>12</sub>H<sub>25</sub>, R' = Me) and 31% VIII (R = C<sub>12</sub>H<sub>25</sub>), m. 107-9°. Cetane (300 ml.) containing 0.01 mole/l. 2-Cl<sub>10</sub>H<sub>7</sub>SH and 0.01 mole II (IV) kept in a 500 ml. open pyrex flask showed rapid peroxidn. followed by a decrease in peroxide content, though the components in cetane showed no peroxidn. After 70 and 23 hrs. the IV-thiol and II-thiol solns. gave some IX, m. 138.5-9.5° (decomposition), and V (R = 2-Cl<sub>10</sub>H<sub>7</sub>, R' = H), m. 145-7°. Cetane (300 ml.) containing 0.3 mole/l. mercaptans and II or IV kept 2 weeks in accelerated storage tests at 43° and the ppts. recrystd. from C<sub>6</sub>H<sub>6</sub>-C<sub>7</sub>H<sub>16</sub> gave the corresponding VIII. Distillation of the mother liquors yielded about 200 mg. 2-indanyl Ph sulfide and 180 mg. 2-indanyl 4-tolyl sulfide, resp. The reported reactions may be important in causing hydrocarbon fuel instability.


IT 102456-32-6, 1-Indanol, 2-(dodecylsulfinyl)-  
(isomer)  
RN 102456-32-6 CAPLUS  
CN 1-Indanol, 2-(dodecylsulfinyl)- (6CI) (CA INDEX NAME)




IT 92621-28-8, 1-Indanol, 2-(phenylsulfinyl)- 93434-18-5,  
1-Indanol, 2-p-tolylsulfinyl- 94305-49-4, 1-Indanol,  
2-(2-naphthylsulfinyl)-  
(isomers)  
RN 92621-28-8 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)



RN 93434-18-5 CAPLUS  
 CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




RN 94305-49-4 CAPLUS  
 CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)




IT 62967-56-0, 1-Indanol, 2-(p-chlorophenylsulfinyl)-  
 92621-28-8, 1-Indanol, 2-(phenylsulfinyl)- 93434-18-5,  
 1-Indanol, 2-p-tolylsulfinyl- 94305-49-4, 1-Indanol,  
 2-(2-naphthylsulfinyl)- 102456-32-6, 1-Indanol,  
 2-(dodecylsulfinyl)-  
 (preparation of)


RN 62967-56-0 CAPLUS  
 CN 1H-Inden-1-ol, 2-[(4-chlorophenyl)sulfinyl]-2,3-dihydro- (9CI) (CA INDEX NAME)




RN 92621-28-8 CAPLUS  
 CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)




RN 93434-18-5 CAPLUS  
CN 1-Indanol, 2-(p-tolylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)



RN 94305-49-4 CAPLUS  
CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)



RN 102456-32-6 CAPLUS  
CN 1-Indanol, 2-(dodecylsulfinyl)- (6CI) (CA INDEX NAME)



L11 ANSWER 20 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 1961:32998 CAPLUS  
DOCUMENT NUMBER: 55:32998  
ORIGINAL REFERENCE NO.: 55:6451b-e  
TITLE: The cooxidation of olefins and mercaptans  
AUTHOR(S): Ford, J. F.; Pitkethly, R. C.; Young, V. O.  
CORPORATE SOURCE: Brit. Petrol. Co. Ltd., Middlesex, UK  
SOURCE: Am. Chem. Soc., Div. Petrol. Chem., Preprints (1957),  
2(No. 1), 111-22  
DOCUMENT TYPE: Journal  
LANGUAGE: Unavailable


AB The primary and secondary products of the cooxidn. of indene (I) and thiophenol (II) were investigated. This was chosen as a simple synthetic system known to be related to the mixts. found in practice (gum-forming). Hydroperoxide was shown to be the main primary product, from which hydroxysulfoxides were formed by spontaneous rearrangement. These results were consistent with those of Kharasch. Results of a typical run at 20° were shown as a graph of O uptake and hydroperoxide and sulfoxide concns. vs. time. Initially almost all the O was used up to form the hydroperoxide (IIa), but concentration of this product fell with increasing sulfoxide concentration IIa was not isolated as such, but was reduced

by LiAlH<sub>4</sub> to trans-2-phenylmercapto-1-indanol (III). The combined reaction mixts. from 3 runs were maintained at 20° until decomposition of the peroxide was complete. From this mixture were isolated and identified by infrared spectrum and mixed m.p. with synthetic compds.




(weight-% given): I, 3.7; trans-2-phenylsulfinyl-1-indanols (IVa), m. 150°, and IVb, m. 99°, total 79.1; III.1.9 and diphenyl disulfide, 1.7; in addition an oxosulfoxide fraction, 8.93 and 7 other fractions in small amts. were separated and characterized by infrared spectra. The following synthetic compds. were prepared for comparison with oxidation products: III and its cis-isomer; IVa and IVb and their cis-isomers of high and low m.p.; cis- and trans-2-phenylsulfonyl-1-indanol; trans-1-phenylmercapto-2-indanol; trans-1-phenylsulfinyl-2-indanols, m. 158° and 134°, resp.; trans-1-phenylsulfonyl-2-indanol, and some intermediates. The synthesis of these compds. was discussed.

IT 92621-28-8, 1-Indanol, 2-(phenylsulfinyl)-  
(stereoisomers)  
RN 92621-28-8 CAPLUS  
CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)



L11 ANSWER 21 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN  
ACCESSION NUMBER: 1960:50305 CAPLUS  
DOCUMENT NUMBER: 54:50305  
ORIGINAL REFERENCE NO.: 54:9855f-i  
TITLE: Organic sulfur compounds. I. Hydroperoxide intermediates in the co-oxidation of mercaptans and olefins  
AUTHOR(S): Oswald, Alexis A.  
CORPORATE SOURCE: Imp. Oil Ltd., Sarnia, Can.  
SOURCE: Journal of Organic Chemistry (1959), 24, 443-4  
CODEN: JOCEAH; ISSN: 0022-3263  
DOCUMENT TYPE: Journal  
LANGUAGE: Unavailable  
AB Air introduced into mixts. of aromatic mercaptans (PhSH, p-MeC6H4SH, p-ClC6H4SH, β-C10H7SH) with reactive olefins (PhCH:CH2, PhCMe:CH2, indene) in hydrocarbons at 0° or lower temps. gave viscous intermediates of high hydroperoxide content. The new S-containing hydroperoxides, RSCHR1CR22OOH usually rearrange at room temperature to the corresponding hydroxyethyl sulfoxides, RSOCHR1CR22OH, with loss of peroxide content, increased absorption in the infrared at 3 and 9-10 μ and by solidification. Indene and β-C10H7SH gave an exceptionally stable co-oxidation product, 2-(2-naphthyl thio)-1-indanyl hydroperoxide, m. 70° (C6H6-C7H16), containing 98% peroxide, rearranging in C6H6 at 40° to give mainly 2-(2-naphthylsulfinyl)-1-indanol, m. 138.5-9.5° (decomposition). Co-oxidation of β-C10H7SH and PhCMe:CH2 gave 2-hydroperoxy-2-phenylpropyl 2-naphthyl sulfide, m. 10°, containing 85% peroxide, rearranged in C6H6 at 40° in 16 hrs. to give 2.3 g./100 ml. sediment, recrystd. (C6H6) to give 2-hydroxy-2-phenylpropyl 2-naphthyl sulfoxide, m. 114-18°. It was suggested that co-oxidation of mercaptans and olefins resulting in formation of hydroperoxides was largely responsible for peroxidation and subsequent color, gum, and sediment formation in untreated petroleum distillates.  
IT 94305-49-4, 1-Indanol, 2-(2-naphthylsulfinyl)-  
(preparation of)  
RN 94305-49-4 CAPLUS  
CN 1-Indanol, 2-(2-naphthylsulfinyl)- (6CI, 7CI) (CA INDEX NAME)



L11 ANSWER 22 OF 22 CAPLUS COPYRIGHT 2004 ACS on STN

ACCESSION NUMBER: 1959:67657 CAPLUS

DOCUMENT NUMBER: 53:67657

ORIGINAL REFERENCE NO.: 53:12253b-i,12254a-i

TITLE: Stereochemistry of the cooxidation products of indene and thiophenol

AUTHOR(S): Ford, J. F.; Pitkethly, R. C.; Young, V. O.

CORPORATE SOURCE: British Petr. Co., Ltd., Sunbury-on-Thames, UK

SOURCE: Tetrahedron (1958), 4, 325-36

CODEN: TETRAB; ISSN: 0040-4020

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB The major primary product of the cooxidation of indene (I) and thiophenol (II) is trans-2-phenylthio-1-indanyl hydroperoxide (III), spontaneously rearranging to the two racemates of trans-2-phenylsulfinyl-1-indanol (IV, V). Attack by the PhS radical on I takes place, within 5%, exclusively at the 2 position and subsequent addition of O to the intermediate 2-phenylthioindanyl radical is, within the same limits, exclusively a trans addition. The cis- and trans-2-phenylthio-1-indanols (VI, VII) and the related sulfoxides and sulfones were synthesized. Under certain conditions the substitution reaction of PhS ion with I bromohydrin is accompanied by migration of the HO group and production of trans-1-phenylthio-2-indanol (VIII). The sulfoxides and sulfones in this series were synthesized. Purification by distillation under N (O content not greater than 0.001%) gave 99.5% pure I, b<sub>13</sub> 67°, n<sub>20D</sub> 1.5765, f.p. - 1.549° and II, b<sub>13</sub> 69°, n<sub>20D</sub> 1.5899. I (0.046 mole) and 0.045 mole II in 150 ml. C<sub>6</sub>H<sub>6</sub> shaken 85 min. at 20 ± 0.1°/760 ± 5 mm. with absorption of 0.043 mole O and the mixture [containing 7 and 77%, resp., of the absorbed O as sulfoxide and hydroperoxide according to the method of Barnard and Hargrave (C.A. 46, 7474i), with addition of 10 ml. AcOH to ensure solubility of the sulfoxides in the aqueous media] kept 140 hrs. at 20° gave a rise in sulfoxide concentration to 45% absorbed O and a complete decay of hydroperoxide according to a 2nd law reaction. The mixture at the end of the oxidation cooled rapidly to 5° and shaken with anhydrous MgSO<sub>4</sub>, the filtered solution added slowly to 4 g. LiAlH<sub>4</sub> in 150 ml. tetrahydrofuran at 0° (hydroperoxide content half-way through the addition, 69% O uptake), the reduction mixture refluxed 6 hrs. and treated with water and aqueous NaOH, extracted with C<sub>6</sub>H<sub>6</sub>, and the product (9.05 g.) chromatographed on silica gel gave 7.19 g. VII (infrared absorption spectra were recorded for all substances described). Separate (3) cooxidations of 5.33 g. I and 4.95 g. II in 300 ml. C<sub>6</sub>H<sub>6</sub> at 20° with 1.347 g. O were carried out, the combined products kept 140 hrs. at 20° to completion of decay of III and deposition of 8.0 g. crystals, m. 140-5°, the solution decanted, and the deposits recrystd. (C<sub>6</sub>H<sub>6</sub>) gave the high-melting isomer IV, m. 150°. The solution diluted with an equal volume of C<sub>5</sub>H<sub>12</sub> and filtered yielded 8.3 g. mixture (of IV and V), m. 80-125°. The filtrate evaporated in vacuo and the residue distilled in a high vacuum to recover 1.3 g. I, the residue triturated with CCl<sub>4</sub>, and the crystalline product (8.3 g.) crystallized (C<sub>6</sub>H<sub>6</sub>-C<sub>6</sub>H<sub>12</sub>) gave the low-melting isomer V, m. 99°. The mother liquors evaporated, the viscous oil (8.76 g.) adsorbed on silica gel, and eluted by solvents of increasing polarity gave 0.6 g. Ph<sub>2</sub>S<sub>2</sub>, m. 61°, 0.65 g. VII, 1.50 g. IV, and 1.4 g. mixture of IV and V, accounting for 86% absorbed O. The

remaining fractions consisted of unidentified sulfides, ketones, oxo sulfides and oxo sulfoxides, as indicated by infrared spectra. IV, V, and the mixture of IV and V oxidized in AcOH with a 5-fold excess of 30% H<sub>2</sub>O<sub>2</sub> gave theoretical yields of trans-2-phenylsulfonyl-1-indanol (IX). A mixture of 95% IX (by weight) and 5% cis-2-phenylsulfonyl-1-indanol (X) began to liquefy at 105° and completely melted at 115°. The crude hydroxy sulfone obtained from the mixture (m. 80-125°) m.

115-17.5° and consequently the % content of cis isomers was less than 5%. I bromohydrin, m. 130° (prepared according to Pope and Read, C.A. 6, 3406), oxidized with CrO<sub>3</sub>-AcOH gave 61% 2-bromo-1-indanone, m. 33.5-4.5°. The bromo ketone (0.2 mole) in 200 ml. MeOH added slowly (N atmospheric) to NaSPh (from 5 g. Na and 0.196 mole II in 150 ml. dry MeOH), the mixture refluxed 15 min., 175 ml. MeOH distilled and the solution refluxed 1 hr., poured into 1 l. water and extracted repeatedly with C<sub>6</sub>H<sub>6</sub>, the washed (5% aqueous NaOH and water) and dried (anhydrous MgSO<sub>4</sub>) extract evaporated, and

the residue recrystd. (C<sub>6</sub>H<sub>12</sub>) gave 48.4 g. product, m. 65-7.5°, recrystd. (C<sub>6</sub>H<sub>6</sub>-C<sub>6</sub>H<sub>12</sub>) to yield pure 2-phenylthio-1-indanone (XI), m. 66-7°; semicarbazone m. 188°; 2,4-dinitrophenylhydrazone m. 191°. XI (4.2 g.) in 30 ml. Et<sub>2</sub>O (dried over Na) added with stirring to 1 g. LiAlH<sub>4</sub> in 50 ml. dry Et<sub>2</sub>O, the mixture refluxed 3 hrs., worked up according to Nystrom and Brown (C.A. 41, 4772g), and the product (4.08 g.) fractionally crystallized (C<sub>6</sub>H<sub>6</sub>-C<sub>6</sub>H<sub>12</sub>) yielded 64% VII, m. 102° (phenylurethan m. 121°), and 14% VI, m. 72°; phenylurethan m. 124°. Desulfurization of VII with Raney Ni gave indan, n<sub>D</sub> 1.5380, identified by its retention volume on a gas chromatography column. Attempted isomerization of VI by heating 4 hrs. with LiAlH<sub>4</sub> in Et<sub>2</sub>O and by refluxing 4 hrs. (N atmospheric) with 10% alc. KOH gave only VI. VII (0.2 g.) refluxed 8 hrs., with 2 ml. 10% alc. KOH, the mixture neutralized with CO<sub>2</sub>, and the filtered and decolorized solution

evaporated

gave 0.12 g. material, m. 80-90°, containing 50% decomposition products other than VI. VII (0.5 g.) heated 4 hrs. in an evacuated sealed tube with 10% alc. KOH at 100° and the mixture worked up with CO<sub>2</sub> gave only unchanged VII. VII (1.15 g.) in AcOH at 5° treated with 10% excess 30% H<sub>2</sub>O<sub>2</sub>, the mixture kept 24 hrs. at room temperature, and the 1.23 g. product fractionally crystallized (C<sub>6</sub>H<sub>6</sub>-C<sub>6</sub>H<sub>12</sub>) gave 0.31 g. IV, m. 149.5-50.5°, and 0.25 g. V, m. 99°. Desulfurization of VII with Raney Ni yielded indan. Similar oxidation of 2.6 g. VI and separation of the colorless gum (2.3 g.) gave the two cis-2-phenylsulfinyl-1-indanols (XII) and (XIII), m. 158°, and 122-3°, resp. The infrared spectra of XII and XIII were similar but differed considerably from those of IV and V. VII (0.75 g.) oxidized in CHCl<sub>3</sub> by shaking with acid aqueous KMnO<sub>4</sub> and working up according to Cunneen (C.A. 41, 3447a) yielded 0.3 g. IX, m. 122°. Oxidation of VI, XII, and XIII in AcOH with 5-fold excess of 30% H<sub>2</sub>O<sub>2</sub> and crystallization of the products from C<sub>6</sub>H<sub>6</sub>-C<sub>6</sub>H<sub>12</sub> yielded

95%

X, m. 131°. Oxidation of VII, IX, and X (2 millimoles) at 40° in 20 min. with 4 millimoles CrO<sub>3</sub> and KMnO<sub>4</sub> oxidation of XI and VI yielded 2-phenylsulfonyl-1-indanone, m. 147-8°; 2,4-dinitrophenylhydrazone m. 262° (EtOAc). The reaction between trans-I bromohydrin (XIV) and NaSPh was found sensitive to reaction conditions and data on quantities of reactants and products formed are tabulated. In expts. 1 and 3, although the concns. were 50% higher in the latter, the proportions were identical (6% excess NaSPh and an addnl. 12% excess PhSH). It is probable that VII and VI were present in both products with the yield of VI appreciably higher in experiment 3 than in

experiment


1. In experiment 2 the excess of NaSPh over XIV was only 1% but the mixture contained NaOEt derived from the 10% excess Na. The product was an oil containing small proportions of VI and VII but the major constituent yielded 2-indanol (XV) on desulfurization with Raney Ni and the reaction must have involved migration of the HO group with formation of VIII. Oxidation of the product (10 g.) in AcOH with 10% excess 30% H<sub>2</sub>O<sub>2</sub>, the oily product (10

g.) triturated with C<sub>6</sub>H<sub>6</sub>, and the solid (4.6 g., m. 145-8°) recrystd. (C<sub>6</sub>H<sub>6</sub>) yielded the high melting isomer of trans-1-phenylsulfinyl-2-indanol (XVI), m. 157-°. The mother liquors evaporated and the residue (4.0 g.) crystallized (1:1 C<sub>6</sub>H<sub>12</sub>-C<sub>6</sub>H<sub>6</sub>) gave the low melting isomer of trans-1-phenylsulfinyl-2-indanol (XVII), m. 133-4.5°. Desulfurization of 0.6 g. XVI in 80 ml. alc. with Raney Ni gave XV, m. 68-9°; phenylurethan m. 131-2°. The final mother liquors gave a solid (0.7 g.), recrystd. (C<sub>6</sub>H<sub>6</sub>-C<sub>6</sub>H<sub>12</sub>) to trans-1-phenylsulfonyl-2-indanol (XVIII), m. 113.5-14.5°. Oxidation of XVI and XVII in AcOH with excess 30% H<sub>2</sub>O<sub>2</sub> gave theoretical yields of XVIII. I epoxide (0.35 mole prepared according to Boeseken and van Loon, C.A. 13, 314) treated according to Bordwell and Kern (C.A. 49, 9639g) with 0.36 mole NaSPh in 50% alc. gave 9.05 g. impure partially dehydrated VIII, n<sub>D</sub> 1.6294; phenylurethan m. 146-7° (C<sub>6</sub>H<sub>12</sub>). By reducing the dielectric constant of the medium and increasing the PhS- concentration the S<sub>N</sub>1 mechanism was inhibited and the S<sub>N</sub>2 promoted. Experiment 4 was carried out with a 4-fold excess of PhS- and PhSH containing a min. amount of alc. was used as solvent. The 89% yield was virtually pure VI, providing convincing confirmation of the assignments of configuration made on the evidence of yields of isomers from the LiAlH<sub>4</sub> reduction of XI, namely that VII, m. 102°, is trans and VI, m. 72°, is cis.

IT 92621-28-8, 1-Indanol, 2-(phenylsulfinyl)-  
(stereoisomers)

RN 92621-28-8 CAPLUS

CN 1H-Inden-1-ol, 2,3-dihydro-2-(phenylsulfinyl)- (9CI) (CA INDEX NAME)



=> file beilstein  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
166.41	542.82

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-15.25	-17.33

CA SUBSCRIBER PRICE

FILE 'BEILSTEIN' ENTERED AT 12:07:55 ON 02 MAY 2004  
COPYRIGHT (c) 2004 Beilstein-Institut zur Foerderung der Chemischen Wissenschaften  
licensed to Beilstein GmbH and MDL Information Systems GmbH

FILE RELOADED ON OCTOBER 20, 2002  
FILE LAST UPDATED ON MARCH 30, 2004

FILE COVERS 1771 TO 2003.  
\*\*\* FILE CONTAINS 8,932,479 SUBSTANCES \*\*\*

>>> PLEASE NOTE: Reaction data and substance data are stored in separate documents and can not be searched together in one query.  
Reaction data for BEILSTEIN compounds may be displayed immediately with the display codes PRE (preparations) and REA (reactions). A substance answer set retrieved after the search for a chemical name, a molecular formula or a structure search

for example can be restricted to compounds with available reaction information by concatenation with PRE/FA, REA/FA or more general with RX/FA. The BEILSTEIN Registry Number (BRN) is the link between a BEILSTEIN compound and belonging reactions. For more detailed reaction searches BRNs can be selected from substance answer sets and searched in the next step as reaction partner BRNs - Reactant (RX.RBRN) or Product BRN (RX.PBRN). After a search for reaction details substance documents associated with reactants or products may be retrieved by searching RX.PBRNs or RX.RBRNs as BRNs. <<<

>>> FOR SEARCHING PREPARATIONS SEE HELP PRE <<<

```
*****
* PLEASE NOTE THAT THERE ARE NO FORMATS FREE OF COST. *
* SET NOTICE FEATURE: THE COST ESTIMATES CALCULATED FOR SET NOTICE *
* ARE BASED ON THE HIGHEST PRICE CATEGORY. THEREFORE; THESE *
* ESTIMATES MAY NOT REFLECT THE ACTUAL COSTS. *
* FOR PRICE INFORMATION SEE HELP COST *
*****
```

=> d his

(FILE 'HOME' ENTERED AT 10:21:40 ON 02 MAY 2004)

FILE 'REGISTRY' ENTERED AT 10:21:48 ON 02 MAY 2004

```
L1      STRUCTURE UPLOADED
L2      1 S L1
L3      8 S L1 FULL
```

FILE 'REGISTRY' ENTERED AT 10:22:48 ON 02 MAY 2004

```
FILE 'CAPLUS' ENTERED AT 10:22:51 ON 02 MAY 2004
L4      3 S L3
```

FILE 'BEILSTEIN' ENTERED AT 11:12:26 ON 02 MAY 2004

```
L5      1 S L1 FULL
```

FILE 'REGISTRY' ENTERED AT 12:00:26 ON 02 MAY 2004

```
L6      STRUCTURE UPLOADED
L7      4 S L6
L8      65 S L6 FULL
L9      30 S L8 NOT SULFONYL
```

FILE 'REGISTRY' ENTERED AT 12:02:30 ON 02 MAY 2004

```
L10     30 S L9
```

FILE 'CAPLUS' ENTERED AT 12:02:37 ON 02 MAY 2004

```
L11     22 S L9
L12     0 S IBIB ABS HITSTR 1-22
```

FILE 'BEILSTEIN' ENTERED AT 12:07:55 ON 02 MAY 2004

=> s l1 full]

MISSING OPERATOR L1 FULL]

The search profile that was entered contains terms or nested terms that are not separated by a logical operator.

=> s l1 full

FULL SEARCH INITIATED 12:10:23 FILE 'BEILSTEIN'  
FULL SCREEN SEARCH COMPLETED - 498 TO ITERATE

100.0% PROCESSED

498 ITERATIONS

1 ANSWERS


SEARCH TIME: 00.00.02

L13 1 SEA SSS FUL L1

=> d ide

L13 ANSWER 1 OF 1 BEILSTEIN COPYRIGHT 2004 BEILSTEIN MDL on STN

Beilstein Records (BRN): 2159196  
Beilstein Pref. RN (BPR): 65495-98-9  
CAS Reg. No. (RN): 65495-98-9  
Chemical Name (CN): 1-Acetoxy-2-p-chlorphenylsulfinylindan  
Autonom Name (AUN): acetic acid 2-(4-chloro-benzenesulfinyl)-  
indan-1-yl ester  
Molec. Formula (MF): C17 H15 Cl O3 S  
Molecular Weight (MW): 334.82  
Lawson Number (LN): 6028, 5223, 1155  
Compound Type (CTYPE): isocyclic  
Constitution ID (CONSID): 1965364  
Tautomer ID (TAUTID): 2106089  
Beilstein Citation (BSO): 5-06  
Entry Date (DED): 1989/06/29  
Update Date (DUPD): 1989/07/26



Field Availability:

Code	Name	Occurrence
BRN	Beilstein Records	1
BPR	Beilstein Preferred RN	1
RN	CAS Registry Number	1
CN	Chemical Name	1
AUN	Autonomname	1
MF	Molecular Formula	1
FW	Formular Weight	1
LN	Lawson Number	3
CTYPE	Compound Type	1
CONSID	Constitution ID	1
TAUTID	Tautomer ID	1
BSO	Beilstein Citation	1
ED	Entry Date	1
UPD	Update Date	1
MP	Melting Point	1

NMR          Nuclear Magnetic Resonance

1

This substance also occurs in Reaction Documents:

Code	Name	Occurrence
=====		
RX	Reaction Documents	1
RXPRO	Substance is Reaction Product	1